Greenyard Crabgrass Preventer with 0.58% Prodiamine

12-0-0

Guaranteed Fertilizer Analysis

Total Nitrogen (N) 12%
0.00% AMMONIACAL NITROGEN
12.00% WATER SOLUBLE NITROGEN *
0.00% WATER INSOLUBLE NITROGEN
Available Phosphate (P_2O_5) 0%
Soluble Potash (K_2O) 0%
Derived from: Urea

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn.
IN MD APPLY AT A MAXIMUM OF .9# OF NITROGEN PER 1,000 SQ. FT.

FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF (excluding golf course putting greens)

STOPS MANY WEED GRASSES BEFORE THEY GROW
PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL

ACTIVE INGREDIENT:

Prodiamine (CAS # 29091-21-2)	0.58%
INERT INGREDIENTS:	99.42%
TOTAL	100.00%

EPA Reg. No. 53883-171-534 EPA Est. No. 75527-PA-12

Keep Out of Reach of Children
CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	• Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician.
IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378.	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (Poa annua); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass[1]; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass3, shepherd's purse2, signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

[1] In many areas a single application of 2.5 to 6 pounds/1,000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds/1,000 square feet (112 to 173 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see"Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

[2] For this weed, apply in late summer, fall or winter prior to germination.

[3] Suppression only.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)

Turf Species	Minimum Rate	Maximum Annual Rate[1]
Bahiagrass, Bermudagrass[2] Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf-Type)	4 lbs/1,000 sq. ft. (175 lbs/acre)	6 lbs/1,000 sq. ft. per year (259 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2 lbs/1,000 sq. ft. (87.5 lbs/acre)	4 lbs/1,000 sq. ft. per year (173 lbs/acre per year)
Creeping Red Fescue	1.5 lbs/1,000 sq. ft. (65.3 lbs/acre)	2.9 lbs/1,000 sq. ft. per year (130 lbs/acre per year)
Creeping Bentgrass	1.5 lbs/1,000 sq. ft. (65.3 lbs/acre)	2.5 lbs/1,000 sq. ft. per year (112 lbs/acre per year)

[1] These are the maximum rates per calendar year.
[2] May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 2 pounds/1,000 square feet (87.5 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map at left.)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
1.9 pounds/1,000 sq. ft. (86 pounds/acre)	4	4	4
2.5 pounds/1,000 sq. ft. (112 pounds/acre)	5	4	4
2.9 pounds/1,000 sq. ft. (130 pounds/acre)	6	5	5
3.16 pounds/1.000 sq. ft. (138 pounds/acre)	-	6	6
4 pounds/1,000 sq. ft. (173 pounds/acre)	-	7	7
4.5 pounds/1,000 sq. ft. (197pounds/acre)	-	-	9
5.14 pounds/1,000 sq. ft. (224 pounds/acre)	-	-	10
5.94 pounds/1,000 sq. ft. (259 pounds/acre)	-	-	12

SUGGESTED SPREADER SETTINGS

Refer to back of bag for spreader settings.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available. If recycling is not available; then dispose of empty bag in a sanitary landfill, or if allowed by State and local authorities, by burning. If burned, say out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Growmark Inc. or the Seller. All such risks shall be assumed by the Buyer.

Growmark Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

GROWMARK INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. GROWMARK INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

Distributed by: Growmark Inc., 1701 Towanda Ave., Bloomington, IL 61701

NET WEIGHT: 50 LBS. (22.6 kg)